



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.


Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

show that, within the limits of ordinary practice, *i. e.*, for angles so small that the difference between arc and sine is inappreciable, the deflection from tangent at  $m$  to set  $n$  is  $(n - m)(n + 2m)$  times the deflection from tangent at 0 to 1.

SOLUTION BY W. J. THOME, Detroit, Michigan.

Let the cubic parabola  $y = ax^3$  be constructed, and on it let the points named 0, 1,  $m$ , and  $n$  be represented by  $O$ ,  $P$ ,  $M$ , and  $N$  respectively. Through these points draw the lines parallel to the axes as shown in the figure. Also the chords  $OP$  and  $MN$ , and  $MT$ , the tangent at  $M$ .



Since the angles are so small that  $\theta = \sin \theta$ , we also have  $\theta = \tan \theta$  and  $\cos \theta = 1$ . Also, an arc of the curve, its chord, and the projection of either on the X-axis are all equal. Hence we have

$$\begin{aligned}\angle LOP &= \tan LOP = \frac{LP}{OL} = \frac{a(OL)^3}{OL} = a(OL)^2 = a(OP)^2 = a(1)^2 = a, \\ \angle SMN &= \tan SMN = \frac{SN}{MS} = \frac{RN - RS}{QR} = \frac{RN - QM}{OR - OQ} = \frac{a(OR)^3 - a(OQ)^3}{OR - OQ} = \frac{a(ON)^3 - a(OM)^3}{ON - OM} \\ &= \frac{a(n)^3 - a(m)^3}{n - m} = \frac{a(n^3 - m^3)}{n - m} = a(n^2 + nm + m^2), \\ \angle SMT &= \tan SMT = \frac{dy}{dx} = 3ax^2 = 3a(OQ)^2 = 3a(OM)^2 = 3am^2, \\ \angle TMN &= \angle SMN - \angle SMT \\ &= a(n^2 + nm + m^2) - 3am^2 = a(n^2 + nm - 2m^2) \\ &= (n - m)(n + 2m)a \\ &= (n - m)(n + 2m) \angle LOP.\end{aligned}$$

**525 (Geometry). Proposed by C. N. SCHMALL, New York City.**

Given a quadrant of a circle  $AOB$ , where  $OA$  and  $OB$  are bounding radii, and a semicircle  $ACO$  having  $OA$  as a diameter and lying on the same side as the quadrant. Describe a circle which shall touch the two arcs and the radius  $OB$ .

**I. SOLUTION BY W. J. THOME, Detroit, Mich.**

Suppose the problem solved and the figure constructed.

Let  $O$  and  $D$  be the centers of the two given circles and let  $E$  be the center of the required circle. Draw  $DE$ . Draw  $OE$  and extend it until it intersects the given quadrant at  $H$ . Through  $E$  draw  $EF$  and  $EG$  perpendicular to  $OA$  and to  $OB$ , respectively.

The problem may be considered solved if we can obtain a value of  $EG$ , the radius of the required circle. Let  $OA = 2R$ ,  $OD = R$ , and  $EG = r$ . Then  $OE = 2R - r$ ,  $DE = R + r$ , and  $DF = R - r$ . Now  $OE^2 - EG^2 = DE^2 - DF^2$ , since  $OG^2 = FE^2$ , or

